## 

## **PERMITTED** GRENA IN ORGANIC SUPERFERRO +S





## **ORGANIC CORRECTIVE - ACTION FERTILIZER NP**



GRENA SUPERFERRO +S is recommended for vineyards, orchards or treating fruit of the forest for iron deficiencies and in alkaline soils

FREE FROM PHOSPHITES AND CHROMIUM VI



**SOURCE** Meatmeal, sulphur and iron sulphate

Physical state: pellet 4 mm

Packaging available: 25 kg bags - 500 kg bags

GRENA SUPERFERRO +S is recommended for the basic fertilisation of
crops that require action of varying degree: iron chlorosis and excessive
limestone are two cases in which it is important to take action with GRENA
SUPERFERRO +S.
the first control of the first

**FARMING** 

It has an excellent corrective action against alkaline soils, balancing the pH; in this way, it helps the plant roots to assimilate the iron humate compounds. This is produced from the interaction between Iron and the GRENA matrix with its humic and fulvic acids and amino acids.

GRENA SUPERFERRO +S ensures a high intake of organic compounds for the general improvement of soil fertility and continuous availability of essential nutrients.

AMINO ACIDS	
Aspartic Acid	1.71 g/100 g
Glutamic Acid	2.71 g/100 g
Alanine	1.16 g/100 g
Arginine	1.21 g/100 g
Phenylalanine	0.83 g/100 g
Glycine	1.71 g/100 g
Hydroxyproline	0.17 g/100 g
Isoleucine	0.83 g/100 g
Histidine	0.34 g/100 g
Leucine	1.58 g/100 g
Lysine	1.00 g/100 g
Proline	1.16 g/100 g
Serine	1.14 g/100 g
Tyrosine	0.64 g/100 g
Threonine	0.89 g/100 g
Valine	1.13 g/100 g
Cysteine and Cystine	0.44 g/100 g
Methionine	1.10 g/100 g
Tryptophan	0.25 g/100 g
FREE AMINO ACIDS	

Glutamic Acid	0.06 g/100 g
Alanine	0.24 g/100 g
Leucine	0.11 g/100 g

MICRO-ELEMENT	S
В	4.62 mg/kg
Fe	661 mg/kg
Mn	37.2 mg/kg
Zn	67.2 mg/kg

COMPOSITION	
Organic matter	60%
Organic substance (SS) (Cx1.724)	54%
Amino acids and proteins (Nx6.25)	18%
Humic and fulvic acids	10%
Humidity	7%
Total nitrogen (N)	3%
Organic nitrogen (N)	3%
Phosphoric anhydride (P <sub>2</sub> O <sub>5</sub> )	2%
Total potassium oxide (K <sub>2</sub> O)	1%
Organic carbon (C)	28%
Sulphuric anhydride (SO <sub>3</sub> )	20%
Total iron (Fe)	3%
Calcium (CaO)	8%
C/N	9.3
рН	< 6

CROP	TIMING*	APPLICATION*	DOSAGE/HA*
Vineyards	autumn - spring	localized distribution per row	600-800 kg/ha
Orchards (pome fruits, stone fruits, citrus fruits etc.)	autumn - spring	localized distribution per row	600-800 kg/ha
Strawberries, soft fruits etc.	pre-sowing or pre-transplant	scatter the product in soil preparation	800-1200 kg/ha
Greenhouse vegetable crops	pre-sowing or pre-transplant	scatter the product in soil preparation	600-800 kg/ha
Open field crops	pre-sowing or pre-transplant	scatter the product in soil preparation	600-800 kg/ha

<sup>\*</sup>guidelines only, for the correct use of our products, please consult a specialist.