

# GRENA BASE

## 3.8.8 S

PERMITTED  
IN ORGANIC  
FARMING



### ORGANO-MINERAL FERTILIZER NPK WITH POTASSIUM SULPHATE




GRENA BASE is recommended for  
basic autumn fertilizations

**FREE FROM  
PHOSPHITES AND  
CHROMIUM VI**

#### SOURCE

Meatmeal  
and feather meal  
Mineral: soft ground  
rock phosphate and  
potassium sulphate



**Physical state:** micro 2 mm - pellet 4 mm

#### Packaging available:

25 kg bags - 500 kg bags

GRENA BASE 3.8.8 is an organo-mineral fertilizer NPK particularly suitable for vineyards and orchards. Through the Grena organic substance conveys nitrogen derived from amino acids of animal origin and from feathermeal. The function of nitrogen is accentuated by the presence of phosphorus (natural phosphorites) allowing the development of robust cell walls. Potassium, in very soluble form and always of biological derivation, allows the formation of sugars in the fruit, culminating in a quality production.

#### AMINO ACIDS IN GRENA MATRIX

Aspartic Acid	1.25 g/100 g
Glutamic Acid	1.62 g/100 g
Alanine	1.02 g/100 g
Arginine	0.83 g/100 g
Phenylalanine	0.56 g/100 g
Glycine	0.95 g/100 g
Hydroxyproline	0.22 g/100 g
Isoleucine	0.62 g/100 g
Histidine	0.31 g/100 g
Leucine	1.10 g/100 g
Lysine	0.56 g/100 g
Proline	0.85 g/100 g
Serine	0.87 g/100 g
Tyrosine	0.33 g/100 g
Threonine	0.59 g/100 g
Valine	0.80 g/100 g
Cysteine and Cystine	0.18 g/100 g
Methionine	0.19 g/100 g
Tryptophan	0.09 g/100 g

#### FREE AMINO ACIDS

Glutamic Acid	0.06 g/100 g
Alanine	0.12 g/100 g
Leucine	0.05 g/100 g

#### MICRO-ELEMENTS

B	2.30 mg/kg
Fe	330 mg/kg
Mn	18.6 mg/kg
Zn	33.6 mg/kg

#### COMPOSITION

Organic matter	40%
<b>Organic substance (SS) (Cx1.724)</b>	<b>36%</b>
Amino acids and proteins (Nx6.25)	20%
Humic and fulvic acids	4%
Humidity	7%
Total nitrogen (N)	3%
<b>Organic nitrogen (N)</b>	<b>3%</b>
Phosphoric anhydride (P <sub>2</sub> O <sub>5</sub> )	8%
<b>Total potassium oxide (K<sub>2</sub>O)</b>	<b>8%</b>
Organic carbon (C)	20%
<b>Calcium (CaO) natural origin</b>	<b>8%</b>
C/N	6.6

CROP	TIMING*	APPLICATION	DOSAGE/HA*
Vineyards	mid-autumn to late spring	localized distribution per row	500-600 kg/ha
Orchards (pome fruits, stone fruits, citrus fruits etc.)	mid-autumn to late spring	localized distribution per row	500-600 kg/ha

\*guidelines only, for the correct use of our products, please consult a specialist.