

GRENA TECH

3.10.5 S +2 MgO

ORGANO-MINERAL FERTILIZER NPK WITH POTASSIUM SULPHATE




GRENA TECH is recommended for soils with phosphorus deficiencies

FREE FROM
PHOSPHITES AND
CHROMIUM VI

SOURCE

Organic: meatmeal and feathermeal
Mineral: soft ground rock phosphate, potassium sulphate and dolomite



Physical state: micro 2 mm - pellet 4 mm

Packaging available:

25 kg bags - 500 kg bags

It is an organo-mineral fertilizer consisting of the union of biological mineral fertilizers with the GRENA organic matrix, rich in proteins, amino acids (naturally biostimulants), humic and fulvic acids.

The levorotatory amino acids in the GRENA organic matrix are the promoters of the development of the secondary roots of the plants and promote the absorption of the nutrients contained directly inside GRENA TECH and those mineralised in the soil.

The naturally present micro-elements are able to catalyse the physiological processes of the plants, allowing to make up for any deficiencies.

The presence of potassium sulphate - in a very soluble form and available for root absorption - promotes the formation of sugars and is therefore essential to ensure high quality productions.

Magnesium provides a considerable input for all vegetable and fruit crops, in order to prevent nutritional deficiencies and ensure the best physiological processes.

AMINO ACIDS IN GRENA MATRIX

| | |
|----------------------|--------------|
| Aspartic Acid | 1.25 g/100 g |
| Glutamic Acid | 1.62 g/100 g |
| Alanine | 1.02 g/100 g |
| Arginine | 0.83 g/100 g |
| Phenylalanine | 0.56 g/100 g |
| Glycine | 0.95 g/100 g |
| Hydroxyproline | 0.22 g/100 g |
| Isoleucine | 0.62 g/100 g |
| Histidine | 0.31 g/100 g |
| Leucine | 1.10 g/100 g |
| Lysine | 0.56 g/100 g |
| Proline | 0.85 g/100 g |
| Serine | 0.87 g/100 g |
| Tyrosine | 0.33 g/100 g |
| Threonine | 0.59 g/100 g |
| Valine | 0.80 g/100 g |
| Cysteine and Cystine | 0.18 g/100 g |
| Methionine | 0.19 g/100 g |
| Tryptophan | 0.09 g/100 g |

FREE AMINO ACIDS

| | |
|---------------|--------------|
| Glutamic Acid | 0.06 g/100 g |
| Alanine | 0.12 g/100 g |
| Leucine | 0.05 g/100 g |

MICRO-ELEMENTS

| | |
|----|------------|
| B | 2.30 mg/kg |
| Fe | 330 mg/kg |
| Mn | 18.6 mg/kg |
| Zn | 33.6 mg/kg |

COMPOSITION

| | |
|---|-----|
| Organic matter | 40% |
| Organic substance (SS) (Cx1.724) | 36% |
| Amino acids and proteins (Nx6.25) | 20% |
| Humic and fulvic acids | 4% |
| Humidity | 7% |
| Total nitrogen (N) | 3% |
| Organic nitrogen (N) | 3% |
| Phosphoric anhydride (P ₂ O ₅) | 10% |
| Total potassium oxide (K ₂ O) | 5% |
| Organic carbon (C) | 20% |
| Sulphuric anhydride (SO ₃) | 6% |
| Magnesium oxide (MgO) of mineral origin | 2% |
| Calcium (CaO) natural origin | 8% |
| C/N | 6.6 |
| pH | 6,5 |

| CROP | TIMING* | APPLICATION* | DOSAGE/HA* |
|--|---------------------------|---|---------------|
| Vineyards | mid-autumn to late spring | localized distribution per row | 500-600 kg/ha |
| Orchards (pome fruits, stone fruits, citrus fruits etc.) | mid-autumn to late spring | localized distribution per row | 500-600 kg/ha |
| Greenhouse vegetable crops | mid-autumn to late spring | scatter the product in soil preparation | 400-500 kg/ha |
| Open field crops (beets) | mid-autumn to late spring | scatter the product in soil preparation | 400-500 kg/ha |
| Flower crops | mid-autumn to late spring | scatter the product in soil preparation | 400-500 kg/ha |

*guidelines only, for the correct use of our products, please consult a specialist.